

Decay Rate Asymmetry of Top Squark

Mayumi Aoki ¹ * and Noriyuki Oshimo ²

¹ *Graduate School of Humanities and Sciences, Ochanomizu University
Otsuka 2-1-1, Bunkyo-ku, Tokyo 112-8610, Japan*

² *Institute for Cosmic Ray Research, University of Tokyo
Midori-cho 3-2-1, Tanashi, Tokyo 188-8502, Japan*

Abstract

We discuss a decay rate asymmetry of the top squark, which is induced by a new source of CP violation intrinsic in the supersymmetric standard model. Although new sources of CP violation in this model are severely constrained from the electric dipole moment of the neutron, an unsuppressed CP -violating phase can still coexist with a top squark whose mass is accessible by near-future colliders. Then the dominant decay mode of the top squark has a width different from its CP conjugate process. The magnitude of this CP asymmetry becomes of order 10^{-3} .

PACS: 11.30.Er, 12.60.Jv, 14.80.Ly

Keywords: CP violation; Supersymmetry; Top squark

*Research Fellow of the Japan Society for the Promotion of Science.

In the standard model (SM) a physical complex phase in quark mass matrices is, aside from the strong CP phase, the only source of CP violation. This phase can explain CP violation in the $K^0\text{-}\bar{K}^0$ system, whereas other sizable CP -violating phenomena are not expected except in the $B^0\text{-}\bar{B}^0$ system [1]. On the other hand, various models beyond the SM contain new sources of CP violation. These effects may be observed in reactions where the SM does not predict CP violation. For instance, near-future collider experiments, such as LHC and NLC, enable precise measurements of certain reactions, which could provide an opportunity of detecting new CP -violating phenomena.

The supersymmetric standard model (SSM) contains a new source of CP violation in the top-squark system [2]. We have recently studied their effects on the decay of the top quark [3]. Taking into account the constraints from the electric dipole moment (EDM) of the neutron, it was shown that a rate asymmetry between the decays $t \rightarrow bW^+$ and $\bar{t} \rightarrow \bar{b}W^-$ can be induced at a magnitude of order 10^{-3} . However, this asymmetry is generated only if there exists a top squark lighter than the top quark. If this condition is not satisfied, other phenomena have to be invoked to search for CP violation arising from the top-squark system. Since CP violation by the top-squark interactions may be able to explain baryon asymmetry of the universe [4], it would be of interest to examine this mechanism of CP violation.

In this letter, we discuss CP violation which is induced by the top squark heavier than the top quark. If a top squark is sufficiently heavy, its dominant decay modes are tree-level two-body decays $\tilde{t} \rightarrow b\omega$ and $\tilde{t} \rightarrow t\chi$, where ω and χ denote a chargino and a neutralino, respectively. These final states are also produced at the one-loop level by final state interactions. If the interactions of the top squark violate CP invariance, these decays are expected to have rates different from their CP conjugate processes, which are measured by the asymmetries

$$A_{CP}^b = \frac{\Gamma(\tilde{t} \rightarrow b\omega^+) - \Gamma(\tilde{t}^* \rightarrow \bar{b}\omega^-)}{\Gamma(\tilde{t} \rightarrow b\omega^+) + \Gamma(\tilde{t}^* \rightarrow \bar{b}\omega^-)}, \quad (1)$$

$$A_{CP}^t = \frac{\Gamma(\tilde{t} \rightarrow t\chi) - \Gamma(\tilde{t}^* \rightarrow \bar{t}\chi)}{\Gamma(\tilde{t} \rightarrow t\chi) + \Gamma(\tilde{t}^* \rightarrow \bar{t}\chi)}. \quad (2)$$

Owing to CPT invariance, the decay widths satisfy the relation

$$\Gamma(\tilde{t} \rightarrow b\omega^+) - \Gamma(\tilde{t}^* \rightarrow \bar{b}\omega^-) = -\left\{\Gamma(\tilde{t} \rightarrow t\chi) - \Gamma(\tilde{t}^* \rightarrow \bar{t}\chi)\right\}, \quad (3)$$

making the total width of the top squark to be the same as that of the anti-top squark. We calculate the asymmetry A_{CP}^b . Its magnitude can become of order 10^{-3} , which may be accessible at near-future colliders.

We first briefly review the model [5]. The SSM contains two new CP -violating phases. These phases can be taken, without loss of generality, for those of the dimensionless coupling constant A in the trilinear terms of the scalar fields and the Higgsino mass parameter m_H in the bilinear term of the Higgs superfields,

$$A = |A| \exp(i\alpha), \quad m_H = |m_H| \exp(i\theta). \quad (4)$$

The mass-squared matrix M_t^2 for the top squarks have these phases, which is given by

$$M_t^2 = \begin{pmatrix} m_t^2 + \cos 2\beta(\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W) M_Z^2 + \tilde{M}_{tL}^2 & m_t(\cot \beta m_H + A^* m_{3/2}) \\ m_t(\cot \beta m_H^* + A m_{3/2}) & m_t^2 + \frac{2}{3} \cos 2\beta \sin^2 \theta_W M_Z^2 + \tilde{M}_{tR}^2 \end{pmatrix}, \quad (5)$$

with $m_{3/2}$ and $\tan \beta$ being respectively the gravitino mass and the ratio of the vacuum expectation values of the Higgs bosons. The mass-squared parameters $\tilde{M}_{tL}^2$ and $\tilde{M}_{tR}^2$ are expressed as

$$\tilde{M}_{tL}^2 = \tilde{M}_q^2 - cm_t^2, \quad \tilde{M}_{tR}^2 = \tilde{M}_q^2 - 2cm_t^2, \quad (6)$$

where $\tilde{M}_q$ is of order $m_{3/2}$ and represents approximately squark masses for the first two generations. The dimensionless constant c stands for an amount of quantum corrections through the Yukawa interactions proportional to the top-quark mass, with $c = 0.1 - 1$.

The mass-squared matrix M_t^2 could become an origin of sizable CP violation, if the imaginary parts of the off-diagonal elements are comparable with the real parts and the diagonal elements. This condition requires that $\tilde{M}_q$ be not much larger than m_t , and at least either θ or α be of order unity. However, these parameters are severely constrained by the EDM of the neutron. An unsuppressed magnitude for θ and a value of $\tilde{M}_q$ smaller than 1 TeV give a too large magnitude to the neutron EDM through one-loop diagrams mediated by charginos and squarks [6]. On the other hand, the experimental upper bound on this EDM is compatible with $\alpha \sim 1$ and $\tilde{M}_q < 1$ TeV, provided that the gluinos are sufficiently heavy [3]. We thus assume that θ is much smaller than unity and α alone has a nonnegligible value.

A nonvanishing value for the asymmetry A_{CP}^b in Eq. (1) is generated, if the decay $\tilde{t} \rightarrow t\chi$, in addition to the decay $\tilde{t} \rightarrow b\omega$, is allowed kinematically. The produced top quark and neutralino can become a bottom quark and a chargino by exchanging charged Higgs bosons $H^\pm$, W bosons, and bottom squarks $\tilde{b}$, as shown in Fig. 1. The interferences of these one-loop diagrams with the tree diagram make the rate

of the decay $\tilde{t} \rightarrow b\omega^+$ different from that of the decay $\tilde{t}^* \rightarrow \bar{b}\omega^-$. The relevant interaction Lagrangian for $\tilde{t}$, $\tilde{b}$, and W is given by

$$\begin{aligned}\mathcal{L} = & i\frac{g}{\sqrt{2}}\tilde{b}_k^\dagger\bar{\omega}_i^c\left(A_{Li}^k\frac{1-\gamma_5}{2}+A_{Ri}^k\frac{1+\gamma_5}{2}\right)t+i\frac{g}{\sqrt{2}}\tilde{t}_k^\dagger\bar{\omega}_i\left(B_{Li}^k\frac{1-\gamma_5}{2}+B_{Ri}^k\frac{1+\gamma_5}{2}\right)b \\ & + i\frac{g}{\sqrt{2}}\tilde{t}_k^\dagger\bar{\chi}_j\left(F_{Lj}^k\frac{1-\gamma_5}{2}+F_{Rj}^k\frac{1+\gamma_5}{2}\right)t+i\frac{g}{\sqrt{2}}\tilde{b}_k^\dagger\bar{\chi}_j\left(G_{Lj}^k\frac{1-\gamma_5}{2}+G_{Rj}^k\frac{1+\gamma_5}{2}\right)b \\ & + \frac{g}{\sqrt{2}}W_\mu^\dagger\bar{\chi}_j\gamma^\mu\left(H_{Lji}\frac{1-\gamma_5}{2}+H_{Rji}\frac{1+\gamma_5}{2}\right)\omega_i+\text{H.c.}\end{aligned}\quad (7)$$

For the definitions of various coefficients, we refer to Ref. [3]. The interaction Lagrangian for $H^\pm$ is given by

$$\begin{aligned}\mathcal{L} = & \frac{g}{\sqrt{2}}H^+\bar{\chi}_j\left(I_{Lji}\frac{1-\gamma_5}{2}+I_{Rji}\frac{1+\gamma_5}{2}\right)\omega_i \\ & + \frac{g}{\sqrt{2}}H^+\bar{t}\left(K_L\frac{1-\gamma_5}{2}+K_R\frac{1+\gamma_5}{2}\right)b+\text{H.c.},\end{aligned}\quad (8)$$

where coefficients are defined as

$$\begin{aligned}I_{Lji} &= -\sin\beta\{(\tan\theta_W N_{1j}+N_{2j})C_{L2i}-\sqrt{2}N_{3j}C_{L1i}\}, \\ I_{Rji} &= \cos\beta\{(\tan\theta_W N_{1j}^*+N_{2j}^*)C_{R2i}+\sqrt{2}N_{4j}^*C_{R1i}\}, \\ K_L &= \cot\beta\frac{m_t}{M_W}, \quad K_R = \tan\beta\frac{m_b}{M_W},\end{aligned}\quad (9)$$

C_L , C_R , and N being the unitary matrices which diagonalize the chargino and neutralino mass matrices [3]. The SSM parameters appearing in our analyses are $\tan\beta$, A , m_H , $\tilde{M}_q$, $m_{3/2}$, c , the SU(2) gaugino mass $\tilde{m}_2$, and the charged Higgs-boson mass $M_{H^\pm}$. Although these parameters are not all independent of each other, they can have various sets of values depending on assumptions for underlying models. For simplicity, we take those parameters independent and assume only rough constraints coming from theoretical and experimental considerations.

We now consider the decay of the lighter top squark into the bottom quark and the lighter chargino $\tilde{t}_1 \rightarrow b\omega_1^+$ and its CP conjugate process. The decay rate asymmetry for these processes are obtained as

$$\begin{aligned}A_{CP}^b = & \frac{\alpha_2}{2}\left[\left\{(|B_{L1}^1|^2+|B_{R1}^1|^2)(\tilde{M}_{t1}^2-m_b^2-\tilde{m}_{\omega1}^2)\right.\right. \\ & \left.\left.-4\text{Re}(B_{L1}^1B_{R1}^{1*})m_b\tilde{m}_{\omega1}\right\}\sqrt{\lambda(\tilde{M}_t^2,\tilde{m}_{\omega1}^2,m_b^2)}\right]^{-1}(T^a+T^b+T^c),\end{aligned}\quad (10)$$

where the kinematic function $\lambda(a, b, c)$ is defined by

$$\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2bc - 2ca. \quad (11)$$

The contributions of the diagrams (a), (b), and (c) in Fig. 1 are represented by T^a , T^b , and T^c , respectively. The terms T^a and T^b are written as

$$T^a = \sum_j \sum_{n=1}^8 \text{Im}(X_{1j1}^n) I_n(\tilde{m}_{\omega 1}, \tilde{m}_{\chi j}, \tilde{M}_{t1}), \quad (12)$$

$$T^b = \sum_j \sum_{n=1}^8 \text{Im}(Y_{1j1}^n) J_n(\tilde{m}_{\omega 1}, \tilde{m}_{\chi j}, \tilde{M}_{t1}), \quad (13)$$

where coefficients X_{1j1}^n , Y_{1j1}^n and functions $I_n(\tilde{m}_{\omega 1}, \tilde{m}_{\chi j}, \tilde{M}_{t1})$, $J_n(\tilde{m}_{\omega 1}, \tilde{m}_{\chi j}, \tilde{M}_{t1})$ are given in Appendix. The term T^c is obtained from T^a by appropriately changing the coefficients and the charged Higgs-boson mass $M_{H\pm}$. The sum for the intermediate neutralinos should be done for those which satisfy the kinematical condition $\tilde{M}_{t1} > m_t + \tilde{m}_{\chi j}$.

In Fig. 2 the absolute value of A_{CP}^b is shown as a function of the SU(2) gaugino mass $\tilde{m}_2$ for $\tilde{M}_q = 400$ GeV and $\alpha = \pi/4$. Four curves correspond to four sets of parameter values for c and $\tan\beta$ listed in Table 1, where the lighter top-squark mass is also given. The other parameters are taken as $|A|m_{3/2} = \tilde{M}_q$, $|m_H| = 100$ GeV, $\theta = 0$, and $M_{H\pm} = 100$ GeV. The parameter $\tilde{m}_2$, together with m_H and $\tan\beta$, determines the masses of the charginos and neutralinos. In the mass ranges where curves are not drawn, the lighter chargino mass becomes smaller than 66 GeV, which is ruled out by LEP2 experiments [7]. In Table 2 the masses of the lighter chargino and the lighter two neutralinos are shown for $\tilde{m}_2 = 500$ GeV. Assuming grand unified models, the gluino mass is also determined by $\tilde{m}_2$. For $\tilde{m}_2 \gtrsim 500$ GeV, the gluinos are heavier than 2 TeV and the value of the neutron EDM lies within its experimental bound.

The asymmetry A_{CP}^b has a magnitude of order 10^{-3} in a region of parameter space where CP violation becomes maximal at the Lagrangian level. For a larger value of $|m_H|$ or $\tilde{M}_q$, $|A_{CP}^b|$ is smaller. The mass of the charged Higgs boson does not affect much the asymmetry. In most region of parameter space in Fig. 2, $\tilde{t}_1$ dominantly decays into $b\omega_1^+$, $t\chi_1$, and $t\chi_2$. The branching ratios of these decays at $\tilde{m}_2 = 500$ GeV are shown in Table 3. For the detection of an asymmetry $A_{CP}^b \sim 10^{-3}$, a necessary number for pairs of $\tilde{t}\tilde{t}^*$ is of order 10^6 . At LHC, the $\tilde{t}\tilde{t}^*$ pairs are expected to be produced at a rate of order 10^5 . If its luminosity is upgraded by one order of magnitude, the asymmetry will be within the reach of detectability.

The interactions which induce the rate asymmetry between the decays $\tilde{t} \rightarrow b\omega^+$ and $\tilde{t}^* \rightarrow \bar{b}\omega^-$ also yield a rate asymmetry between the decays $\tilde{t} \rightarrow t\chi$ and $\tilde{t}^* \rightarrow \bar{t}\chi$, satisfying the relation in Eq. (3). As seen in Table 3, the width of $\tilde{t}_1 \rightarrow t\chi_1$ or

$\tilde{t}_1 \rightarrow t\chi_2$ is generally several times smaller than that of $\tilde{t}_1 \rightarrow b\omega_1^+$, and accordingly the former decay rate asymmetry becomes larger than the latter by the same order of magnitude. It will also be possible to examine CP violation through the decay into $t\chi_1$ or $t\chi_2$. However, the relation in Eq. (3) makes the detection of a decay rate asymmetry a little involved. The top quark decays into b and W , while the chargino decays through χ_1 and W^* , where W^* denotes the virtual state of the W boson. Consequently, the top-squark decays $\tilde{t}_1 \rightarrow b\omega_1^+$ and $\tilde{t}_1 \rightarrow t\chi_1$ have the same particles in their final states. In order to measure the asymmetry, these decays have to be distinguished. Such a distinction can be made by e.g. examining the energy of the b quark. On the other hand, the decay $\tilde{t}_1 \rightarrow t\chi_2$ shows a different topology, because of the subsequent decay of the second lightest neutralino χ_2 .

In summary, we have studied the decay rate asymmetry of the lighter top squark. By this asymmetry, CP violation intrinsic in the SSM can be probed. If the gluinos have a mass larger than 1 TeV, a new CP -violating phase contained in the top-squark mass-squared matrix is not constrained much from the EDM of the neutron. For a top-squark mass around 300 GeV with the CP -violating phase of order unity, the asymmetry is of order 10^{-3} , which may be detectable in the near future.

The work of M.A. is supported in part by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture, Japan.

Appendix

The coefficients X_{ijk}^n and the functions $I_n(\tilde{m}_\omega, \tilde{m}_\chi, \tilde{M}_t)$ in Eq. (12) are defined by

$$\begin{aligned}
X_{ijk}^1 &= K_L^* B_{Li}^k F_{Lj}^{k*} I_{Lji} + K_R^* B_{Ri}^k F_{Rj}^{k*} I_{Rji}, \\
X_{ijk}^2 &= K_L^* B_{Li}^k F_{Rj}^{k*} I_{Lji} + K_R^* B_{Ri}^k F_{Lj}^{k*} I_{Rji}, \\
X_{ijk}^3 &= K_L^* B_{Ri}^k F_{Lj}^{k*} I_{Lji} + K_R^* B_{Li}^k F_{Rj}^{k*} I_{Rji}, \\
X_{ijk}^4 &= K_L^* B_{Ri}^k F_{Rj}^{k*} I_{Lji} + K_R^* B_{Li}^k F_{Lj}^{k*} I_{Rji}, \\
X_{ijk}^5 &= K_R^* B_{Li}^k F_{Lj}^{k*} I_{Lji} + K_L^* B_{Ri}^k F_{Rj}^{k*} I_{Rji}, \\
X_{ijk}^6 &= K_R^* B_{Li}^k F_{Rj}^{k*} I_{Lji} + K_L^* B_{Ri}^k F_{Lj}^{k*} I_{Rji}, \\
X_{ijk}^7 &= K_R^* B_{Ri}^k F_{Lj}^{k*} I_{Lji} + K_L^* B_{Li}^k F_{Rj}^{k*} I_{Rji}, \\
X_{ijk}^8 &= K_R^* B_{Ri}^k F_{Rj}^{k*} I_{Lji} + K_L^* B_{Li}^k F_{Lj}^{k*} I_{Rji},
\end{aligned}$$

$$\begin{aligned}
I_1 &= \frac{1}{2} \tilde{m}_\omega m_t \left\{ T + (m_t^2 + \tilde{m}_\omega^2 - \tilde{M}_t^2 - M_{H^\pm}^2) \log \left| \frac{S+T}{S-T} \right| \right\}, \\
I_2 &= \frac{1}{2} \tilde{m}_\omega \tilde{m}_\chi \left\{ T + (m_t^2 + m_b^2 - M_{H^\pm}^2) \log \left| \frac{S+T}{S-T} \right| \right\}, \\
I_3 &= \frac{1}{2} m_b m_t \left\{ T + (\tilde{m}_\chi^2 + \tilde{m}_\omega^2 - M_{H^\pm}^2) \log \left| \frac{S+T}{S-T} \right| \right\}, \\
I_4 &= \frac{1}{2} m_b \tilde{m}_\chi \left\{ T + (\tilde{m}_\chi^2 + m_b^2 - \tilde{M}_t^2 - M_{H^\pm}^2) \log \left| \frac{S+T}{S-T} \right| \right\}, \\
I_5 &= -\frac{1}{2} \tilde{m}_\omega m_b (\tilde{M}_t^2 - m_t^2 - \tilde{m}_\chi^2) \log \left| \frac{S+T}{S-T} \right|, \\
I_6 &= m_b m_t \tilde{m}_\omega \tilde{m}_\chi \log \left| \frac{S+T}{S-T} \right|, \\
I_7 &= \frac{1}{2} \left\{ \tilde{M}_t^2 T + (-\tilde{M}_t^2 M_{H^\pm}^2 + \tilde{m}_\omega^2 m_t^2 + m_b^2 \tilde{m}_\chi^2) \log \left| \frac{S+T}{S-T} \right| \right\}, \\
I_8 &= -\frac{1}{2} \tilde{m}_\chi m_t (\tilde{M}_t^2 - m_b^2 - \tilde{m}_\omega^2) \log \left| \frac{S+T}{S-T} \right|,
\end{aligned}$$

where

$$\begin{aligned}
S &= \frac{1}{\tilde{M}_t^2} (\tilde{M}_t^2 + \tilde{m}_\omega^2 - m_b^2) (\tilde{M}_t^2 + \tilde{m}_\chi^2 - m_t^2) - 2(\tilde{m}_\chi^2 + \tilde{m}_\omega^2 - M_{H^\pm}^2), \\
T &= \frac{1}{\tilde{M}_t^2} \sqrt{\lambda(m_b^2, \tilde{M}_t^2, \tilde{m}_\omega^2) \lambda(m_t^2, \tilde{M}_t^2, \tilde{m}_\chi^2)}.
\end{aligned}$$

The coefficients Y_{ijk}^n and the functions $J_n(\tilde{m}_\omega, \tilde{m}_\chi, \tilde{M}_t)$ in Eq. (13) are defined by

$$\begin{aligned}
Y_{ijk}^1 &= B_{Li}^k F_{Lj}^{k*} H_{Lji}, & Y_{ijk}^2 &= B_{Li}^k F_{Lj}^{k*} H_{Rji}, & Y_{ijk}^3 &= B_{Li}^k F_{Rj}^{k*} H_{Lji}, \\
Y_{ijk}^4 &= B_{Li}^k F_{Rj}^{k*} H_{Rji}, & Y_{ijk}^5 &= B_{Ri}^k F_{Lj}^{k*} H_{Lji}, & Y_{ijk}^6 &= B_{Ri}^k F_{Lj}^{k*} H_{Rji}, \\
Y_{ijk}^7 &= B_{Ri}^k F_{Rj}^{k*} H_{Lji}, & Y_{ijk}^8 &= B_{Ri}^k F_{Rj}^{k*} H_{Rji}, \\
J_1 &= \frac{1}{2} \tilde{m}_\omega \tilde{m}_\chi \left[\frac{2M_W^2 + m_t^2 + m_b^2}{M_W^2} T \right. \\
&\quad \left. + \left\{ m_b^2 + m_t^2 - 2M_W^2 + \frac{(m_b^2 - m_t^2)^2}{M_W^2} \right\} \log \left| \frac{S'+T}{S'-T} \right| \right], \\
J_2 &= -\frac{m_b^2 \tilde{m}_\chi^2 + m_t^2 \tilde{m}_\omega^2}{2M_W^2} T - \frac{1}{2} \left[2(\tilde{M}_t^2 - \tilde{m}_\chi^2 - m_t^2)(\tilde{M}_t^2 - \tilde{m}_\omega^2 - m_b^2) \right. \\
&\quad \left. - m_b^2 \tilde{m}_\chi^2 - m_t^2 \tilde{m}_\omega^2 + \frac{1}{M_W^2} \left\{ (m_b^2 \tilde{m}_\chi^2 - m_t^2 \tilde{m}_\omega^2)(\tilde{m}_\chi^2 - \tilde{m}_\omega^2 + m_b^2 - m_t^2) \right. \right. \\
&\quad \left. \left. + (m_b^2 - m_t^2)(\tilde{m}_\omega^2 - \tilde{m}_\chi^2) \tilde{M}_t^2 \right\} \right] \log \left| \frac{S'+T}{S'-T} \right|,
\end{aligned}$$

$$\begin{aligned}
J_3 &= \frac{1}{2} \tilde{m}_\omega m_t \left[\frac{2M_W^2 - \tilde{M}_t^2 + \tilde{m}_\chi^2 + m_b^2}{M_W^2} T + \left\{ 2(\tilde{m}_\omega^2 - M_W^2 + m_t^2) \right. \right. \\
&\quad \left. \left. - \tilde{M}_t^2 - m_b^2 - \tilde{m}_\chi^2 + \frac{(m_b^2 - m_t^2)(\tilde{m}_\omega^2 - \tilde{m}_\chi^2)}{M_W^2} \right\} \log \left| \frac{S' + T}{S' - T} \right| \right], \\
J_4 &= \frac{\tilde{m}_\chi m_t (\tilde{M}_t^2 - m_b^2 - \tilde{m}_\omega^2)}{2M_W^2} \left(T + 3M_W^2 \log \left| \frac{S' + T}{S' - T} \right| \right), \\
J_5 &= \frac{1}{2} \tilde{m}_\chi m_b \left[\frac{2M_W^2 - \tilde{M}_t^2 + \tilde{m}_\omega^2 + m_t^2}{M_W^2} T + \left\{ 2(\tilde{m}_\chi^2 - M_W^2 + m_b^2) \right. \right. \\
&\quad \left. \left. - \tilde{M}_t^2 - m_t^2 - \tilde{m}_\omega^2 + \frac{(m_b^2 - m_t^2)(\tilde{m}_\omega^2 - \tilde{m}_\chi^2)}{M_W^2} \right\} \log \left| \frac{S' + T}{S' - T} \right| \right], \\
J_6 &= \frac{\tilde{m}_\omega m_b (\tilde{M}_t^2 - m_t^2 - \tilde{m}_\chi^2)}{2M_W^2} \left(T + 3M_W^2 \log \left| \frac{S' + T}{S' - T} \right| \right), \\
J_7 &= \frac{1}{2} m_t m_b \left[\frac{2M_W^2 + \tilde{m}_\chi^2 + \tilde{m}_\omega^2}{M_W^2} T \right. \\
&\quad \left. + \left\{ \tilde{m}_\chi^2 + \tilde{m}_\omega^2 - 2M_W^2 + \frac{(\tilde{m}_\chi^2 - \tilde{m}_\omega^2)^2}{M_W^2} \right\} \log \left| \frac{S' + T}{S' - T} \right| \right], \\
J_8 &= -\frac{m_b m_t \tilde{m}_\omega \tilde{m}_\chi}{M_W^2} \left(T + 3M_W^2 \log \left| \frac{S' + T}{S' - T} \right| \right),
\end{aligned}$$

where S' is derived from S by changing $M_{H^\pm}$ to M_W .

References

- [1] For general reviews on CP violation, see e.g. J.F. Donoghue, B.R. Holstein, and G. Valencia, Int. J. Mod. Phys. A2 (1987) 319;
W. Grimus, Fortschr. Phys. 36 (1988) 201.
- [2] C. R. Schmidt, Phys. Lett. B 293 (1992) 111;
E. Christova and M. Fabbrihesi, Phys. Lett. B315 (1993) 338; B320 (1994) 299;
B. Grzadkowski and W.-Y. Keung, Phys. Lett. B316 (1993) 137; B319 (1993) 526;
A. Bartl, E. Christova, and W. Majerotto, Nucl. Phys. B460 (1996) 235;
A. Bartl, E. Christova, T. Gajdosik, and W. Majerotto, Nucl. Phys. B507 (1997) 35;
S. Bar-Shalom, D. Atwood, and A. Soni, Phys. Rev. D57 (1998) 1495.
- [3] M. Aoki and N. Oshimo, hep-ph/9801294 (to appear in Nucl. Phys. B).
- [4] M. Aoki, A. Sugamoto, and N. Oshimo, Prog. Theor. Phys. 98 (1997) 1325.
- [5] See e.g. H.P. Nilles, Phys. Rep. 110 (1984) 1;
P. Nath, R. Arnowitt, and A.H. Chamseddine, *Applied N=1 Supergravity* (World Scientific, Singapore, 1984);
H.E. Haber and G.L. Kane, Phys. Rep. 117 (1985) 75.
- [6] Y. Kizukuri and N. Oshimo, Phys. Rev. D45 (1992) 1806; D46 (1992) 3025.
- [7] Particle Data Group, Eur. Phys. J. C3 (1998) 1.

Table 1: The parameter values for curves (i)–(iv) in Fig. 2. The lighter top-squark mass is also given.

	c	$\tan \beta$	$\tilde{M}_{t1}$ (GeV)
(i)	0.1	2	328
(ii)	0.1	10	335
(iii)	0.5	2	296
(iv)	0.5	10	304

Table 2: The chargino and neutralino masses at $\tilde{m}_2 = 500$ GeV for the parameter values of curves (i)–(iv) in Fig. 2.

	$\tilde{m}_{\omega 1}$	$\tilde{m}_{\chi 1}$	$\tilde{m}_{\chi 2}$	(GeV)
(i),(iii)	87	77	102	
(ii),(iv)	95	85	107	

Table 3: The branching ratios for the four sets of parameter values in Table 1 with $\tilde{m}_2 = 500$ GeV.

	$\tilde{t}_1 \rightarrow b\omega_1^+$	$\tilde{t}_1 \rightarrow t\chi_1$	$\tilde{t}_1 \rightarrow t\chi_2$
(i)	0.59	0.14	0.27
(ii)	0.59	0.13	0.28
(iii)	0.72	0.09	0.19
(iv)	0.71	0.09	0.20

Figure 1: The one-loop diagrams for the decay of a top squark into a bottom quark and a chargino.




Figure 2: The decay rate asymmetry as a function of $\tilde{m}_2$. Four curves correspond to four sets of values for c and $\tan \beta$ in Table 1.

